arXiv:1212.4681v1 [math.CA] 5 Dec 2012

Inequalities for Generalized Trigonometric and Hyperbolic Sine Functions

Miao-Kun Wang[1], Yu-Ming Chu[2,*] and Yue-Ping Jiang[3]

[1]Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China;
[2]School of Mathematics and Computation Sciences, Hunan City University, Yiyang 413000, China;
[3]College of Mathematics and Econometrics, Hunan University, Changsha 410082, China.
Correspondence should be addressed to Yu-Ming Chu, chuyuming@hutc.zj.cn

Abstract: We prove that the inequalities $\sin_{p,q}(\sqrt{rs}) \geq \sqrt{\sin_{p,q}(r)\sin_{p,q}(s)}$ and $\sinh_{p,q}(\sqrt{r^*s^*}) \leq \sqrt{\sinh_{p,q}(r^*)\sinh_{p,q}(s^*)}$ hold for all $p,q \in (1,\infty)$, $r,s \in (0, \int_0^1 (1-t^q)^{-1/p}dt)$ and $r^*, s^* \in (0, \int_0^\infty (1+t^q)^{-1/p}dt)$, where $\sin_{p,q}$ and $\sinh_{p,q}$ are the generalized trigonometric and hyperbolic sine functions, respectively. As a consequence of the results, we prove a conjecture due to Bhayo and Vuorinen [J. Approx. Theory, 164(2012)].

Keywords: generalized trigonometric function, generalized hyperbolic function, inequality
2010 Mathematics Subject Classification: 33B10

1. Introduction

It is well known from basic calculus that

$$\frac{\pi}{2} = \int_0^1 \frac{1}{\sqrt{1-t^2}}dx$$

and

$$\arcsin(x) = \int_0^x \frac{1}{\sqrt{1-t^2}}dt, \quad 0 \leq x \leq 1.$$

Since the function $\arcsin(x)$ is a differentiable function on $[0,1]$ and $t \to 1/\sqrt{1-t^2}$ is strictly increasing on $[0,1)$, we can define sin on $[0,\pi/2]$ as the inverse function of arcsin. By standard extension procedures we can define the sin function on $(-\infty,\infty)$.

For $p,q > 1$, let

$$F_{p,q}(x) = \int_0^x (1-t^q)^{-1/p}dt, \quad x \in [0,1],$$

$$\frac{\pi_{p,q}}{2} = \int_0^1 (1-t^q)^{-1/p} dt.$$

Then $F_{p,q} : [0,1] \to [0, \pi_{p,q}/2]$ is an increasing homeomorphism, denoted by $\mathrm{arcsin}_{p,q}$. Thus its inverse

$$\sin_{p,q} = F_{p,q}^{-1}$$

is defined on the interval $[0, \pi_{p,q}/2]$. By the similar extension as the sine function, we can get a differentiable function $\sin_{p,q}$ defined on $\mathbb{R}$. We call $\sin_{p,q}$ the generalized (p,q)-trigonometric sine function.

We also defined $\mathrm{arccos}_{p,q}(x) = \mathrm{arcsin}_{p,q}((1-x^p)^{1/q})$ (See [2, 7]), and the inverse of generalized (p,q)-hyperbolic sine function

$$\mathrm{arcsinh}_{p,q}(x) = \int_0^x (1+t^q)^{-1/p} dt, \quad x \in (0,\infty).$$

Their inverse functions are

$$\sin_{p,q} : (0, \pi_{p,q}/2) \to (0,1), \qquad \cos_{p,q} : (0, \pi_{p,q}/2) \to (0,1),$$
$$\sinh_{p,q} : (0, m^*_{p,q}) \to (0,\infty), \qquad m^*_{p,q} = \int_0^\infty (1+t^q)^{-1/p} dt.$$

When $p = q$, the (p,q)-functions $\sin_{p,q}$, $\cos_{p,q}$, $\sinh_{p,q}$, $\mathrm{arcsin}_{p,q}$, $\mathrm{arccos}_{p,q}$ and $\mathrm{arcsinh}_{p,q}$ reduce to p-functions $\sin_p$, $\cos_p$, $\sinh_p$, arcsin_p, arccos_p and $\mathrm{arcsinh}_p$ (See [5, 8, 10]), respectively. In particular, when $p = q = 2$, the (p,q)-functions become our familiar trigonometric and hyperbolic functions.

Recently, the generalized trigonometric and hyperbolic functions ((p,q)-functions and p-functions) have been found many important applications in differential equations, the theory of operator, approximation theory and other related fields [4, 9, 11].

In face of the importance of generalized trigonometric and hyperbolic functions they have been studied by many authors from different points of view [1, 3-8, 10, 12]. Ednumds, Gurka and Lang [7] gave the basic properties of generalized (p,q)-trigonometric functions, and proved that

$$\sin_{4/3,4}(2x) = \frac{2\sin_{4/3,4}(x)(\cos_{4/3,4}(x))^{1/3}}{(1+4(\sin_{4/3,4}(x))^4(\cos_{4/3,4}(x))^{4/3})^{1/2}}.$$

Klén, Vuorinen and Zhang [8] generalized some classical inequalities for trigonometric and hyperbolic functions, such as Mitrinović-Adamović inequality and Lazarević's inequality.

Bhayo and Vuorinen [2] found that the functions $\mathrm{arcsin}_{p,q}$ and $\mathrm{arcsinh}_{p,q}$ can be expressed in terms of Gaussian hypergeometric functions. Applying the vast available information about the hypergeometric functions, some remarkable properties and inequalities for generalized trigonometric and hyperbolic functions are obtained. Moreover, they raised the following conjecture.

Conjecture 1.1. If $p, q \in (1,\infty)$ and $r, s \in (0,1)$, then

$$\sin_{p,q}(\sqrt{rs}) \geq \sqrt{\sin_{p,q}(r)\sin_{p,q}(s)},$$

$$\sinh_{p,q}(\sqrt{rs}) \leq \sqrt{\sinh_{p,q}(r)\sinh_{p,q}(s)}.$$

The main purpose of this paper is to give a positive answer to the Conjecture 1.1 and generalized the inequalities in Conjecture 1.1. Our main result is the following Theorem 1.1.

Theorem 1.1. If $p, q \in (1, \infty)$, then
(1) Inequality

$$\sin_{p,q}(\sqrt{rs}) \geq \sqrt{\sin_{p,q}(r)\sin_{p,q}(s)} \tag{1.1}$$

holds for all $r, s \in (0, \pi_{p,q}/2)$.
(2) Inequality

$$\sinh_{p,q}(\sqrt{r^* s^*}) \leq \sqrt{\sinh_{p,q}(r^*)\sinh_{p,q}(s^*)} \tag{1.2}$$

holds for all $r^*, s^* \in (0, m^*_{p,q})$.

2. Proof of Theorem 1.1

In order to prove Theorem 1.1 and Conjecture 1.1, we present three Lemmas at first.

Lemma 2.1. If $p, q \in (1, \infty)$, then inequality

$$\text{arcsin}_{p,q}(x) > \frac{px(1-x^q)^{1-1/p}}{(q-p)x^q + p} \tag{2.1}$$

holds for $x \in (0, 1)$.

Proof. Let

$$\zeta(x) = \text{arcsin}_{p,q}(x) - \frac{px(1-x^q)^{1-1/q}}{(q-p)x^q + p}, \quad x \in (0, 1). \tag{2.2}$$

Then simple computations lead to

$$\zeta(0) = 0, \tag{2.3}$$

$$\begin{aligned}\zeta'(x) =& \frac{1}{(1-x^q)^{1/p}} - \frac{p^2 + p(2q - 2p - q^2)x^q + (q-p)^2 x^{2q}}{(1-x^q)^{1/p}(p - px^q + qx^q)^2} \\ =& \frac{pq^2 x^q}{(1-x^q)^{1/p}(p - px^q + qx^q)^2} > 0 \end{aligned} \tag{2.4}$$

for $x \in (0, 1)$.
Therefore, Lemma 2.1 follows easily from (2.2)-(2.4).

Lemma 2.2. If $p, q \in (1, \infty)$, then inequality

$$\frac{x}{\operatorname{arcsinh}_{p,q}(x)} > \frac{(p-q)x^q + p}{p(1+x^q)^{1-1/p}} \tag{2.5}$$

holds for $x \in (0, \infty)$.

Proof. We divide the proof into two cases.
Case 1 $p \geq q$. Let

$$\eta(x) = \operatorname{arcsinh}_{p,q}(x) - \frac{px(1+x^q)^{1-1/p}}{(p-q)x^q + p}, \quad x \in (0, \infty), \tag{2.6}$$

then simple computations lead to

$$\eta(0) = 0, \tag{2.7}$$

$$\begin{aligned} \eta'(x) =& \frac{1}{(1+x^q)^{1/p}} - \frac{p^2 + p(2p - 2q + q^2)x^q + (q-p)^2 x^{2q}}{(1+x^q)^{1/p}(p - qx^q + px^q)^2} \\ =& - \frac{pq^2 x^q}{(1+x^q)^{1/p}(p - qx^q + px^q)^2} < 0 \end{aligned} \tag{2.8}$$

for $x \in (0, \infty)$.
Therefore, inequality (2.5) follows easily from (2.6)-(2.8) and $p \geq q$.

Case 2 $p < q$. Let $x_0 = [p/(q-p)]^{1/q}$, then $(p-q)x^q + p > 0$ for $x \in (0, x_0)$ and $(p-q)x^q + p < 0$ for $x \in (x_0, \infty)$. Thus it is sufficient to prove inequality (2.5) for $x \in (0, x_0)$, which easily follows from the proof of Case 1.

Lemma 2.3. If $p, q > 1$, then $m^*_{p,q} = \int\limits_0^\infty (1+t^q)^{-1/p} dt > 1$.

Proof. From the basic properties of generalized integrals we clearly see that $m^*_{p,q} = +\infty$ if $p \geq q$ and $m^*_{p,q} < +\infty$ if $p < q$. Since

$$(1 + t^{(q-2p)/p})^p > 1 + t^{q-2p} > 1 + t^q$$

for $t \in (0, 1)$, we get

$$\begin{aligned} m^*_{p,q} =& \int_0^\infty \frac{1}{(1+t^q)^{1/p}} dt = \int_0^1 \frac{1}{(1+t^q)^{1/p}} dt + \int_1^\infty \frac{1}{(1+t^q)^{1/p}} dt \\ =& \int_0^1 \frac{1}{(1+t^q)^{1/p}} dt + \int_0^1 \frac{t^{q/p-2}}{(1+t^q)^{1/p}} dt = \int_0^1 \frac{1 + t^{q/p-2}}{(1+t^q)^{1/p}} dt > 1. \end{aligned}$$

Proof of Theorem 1.1. For part (1), without loss of generality, we assume that $0 < x \leq y < 1$. Define

$$J(x, y) = \frac{\arcsin_{p,q}(H_{p_1}(x, y))^2}{\arcsin_{p,q}(x) \arcsin_{p,q}(y)}, \quad p_1 \in \mathbb{R}, \tag{2.9}$$

where

$$H_{p^*}(a,b)=\begin{cases}\left(\frac{a^{p^*}+b^{p^*}}{2}\right)^{1/p^*}, & p^*\neq 0,\\ \sqrt{ab}, & p^*=0\end{cases}$$

is the Hölder mean of order $p^*\in\mathbb{R}$ of two positive numbers a and b.

Let $t=H_{p_1}(x,y)$, then $\partial t/\partial x=(x/t)^{p_1-1}/2$. If $x<y$, then $t>x$. By logarithmic differentiation, we get

$$\frac{1}{J(x,y)}\frac{\partial J}{\partial x}=x^{p_1-1}\left[\frac{t^{1-p_1}}{\operatorname{arcsin}_{p,q}(t)(1-t^q)^{1/p}}-\frac{x^{1-p_1}}{\operatorname{arcsin}_{p,q}(x)(1-x^q)^{1/p}}\right]. \tag{2.10}$$

Let

$$F(x)=\frac{x^{1-p_1}}{\operatorname{arcsin}_{p,q}(x)(1-x^q)^{1/p}},\quad x\in(0,1), \tag{2.11}$$

then logarithmic differentiation yields

$$\begin{aligned}\frac{F'(x)}{F(x)}=&(1-p_1)\frac{1}{x}-\frac{1}{\operatorname{arcsin}_{p,q}(x)(1-x^q)^{1/p}}+\frac{qx^{q-1}}{p(1-x^q)}\\ =&\frac{1}{x}(G(x)+1-p_1),\end{aligned} \tag{2.12}$$

where

$$G(x)=\frac{qx^q}{p(1-x^q)}-\frac{x}{\operatorname{arcsin}_{p,q}(x)(1-x^q)^{1/p}}.$$

Note that

$$\lim_{x\to 0}G(x)=-1,\quad \lim_{x\to 1}G(x)=+\infty. \tag{2.13}$$

It follows from Lemma 2.1 and (2.13) that the range of $G(x)$ is $(-1,\infty)$. Thus from (2.12) we conclude that $F'(x)\geq 0$ for $x\in(0,1)$ if and only if $p_1\leq 0$. Namely, $F(x)$ is strictly increasing on $(0,1)$ if and only if $p_1\leq 0$. Moreover, if $p_1>0$, then $F(x)$ is not monotone.

Next, we divide the proof into two cases.

Case A $p_1\leq 0$. Then from equation (2.10) and the monotonicity of $F(x)$ we clearly see that $\partial J/\partial x>0$. Hence $J(x,y)\leq J(y,y)=1$. Then by (2.9) we get

$$\operatorname{arcsin}_{p,q}(H_{p_1}(x,y))\leq\sqrt{\operatorname{arcsin}_{p,q}(x)\operatorname{arcsin}_{p,q}(y)} \tag{2.14}$$

for $p_1\leq 0$, with equality if and only if $x=y$.

Case B $p_1>0$. Then using the similar argument in Case A, we conclude that there exists $x_1,x_2,y_1,y_2\in(0,1)$ such that

$$\operatorname{arcsin}_{p,q}(H_{p_1}(x_1,y_1))<\sqrt{\operatorname{arcsin}_{p,q}(x_1)\operatorname{arcsin}_{p,q}(y_1)},$$

$$\operatorname{arcsin}_{p,q}(H_{p_1}(x_2,y_2))>\sqrt{\operatorname{arcsin}_{p,q}(x_2)\operatorname{arcsin}_{p,q}(y_2)}.$$

Finally, taking $x = \sin_{p,q}(r)$ and $y = \sin_{p,q}(s)$ in inequality (2.14), we have

$$\sin_{p,q}(\sqrt{rs}) \geq H_{p_1}(\sin_{p,q}(r), \sin_{p,q}(s)) \tag{2.15}$$

for all $r, s \in (0, \pi_{p,q}/2)$ if and only if $p_1 \leq 0$. In particular, inequality (1.1) follows from (2.15) with $p_1 = 0$.

For part (2), without loss of generality, we assume that $0 < x \leq y < \infty$. Define

$$J^*(x,y) = \frac{\operatorname{arcsinh}_{p,q}(H_{p_2}(x,y))^2}{\operatorname{arcsinh}_{p,q}(x)\operatorname{arcsinh}_{p,q}(y)}, \quad p_2 \in \mathbb{R}. \tag{2.16}$$

Let $t = H_{p_2}(x,y)$, then $\partial t/\partial x = (x/t)^{p_2-1}/2$. If $x < y$, then $t > x$. By logarithmic differentiation, we get

$$\frac{1}{J^*(x,y)}\frac{\partial J^*}{\partial x} = x^{p_2-1}\left[\frac{t^{1-p_2}}{\operatorname{arcsinh}_{p,q}(t)(1+t^q)^{1/p}} - \frac{x^{1-p_2}}{\operatorname{arcsinh}_{p,q}(x)(1+x^q)^{1/p}}\right]. \tag{2.17}$$

Let

$$F^*(x) = \frac{x^{1-p_2}}{\arcsin_{p,q}(x)(1+x^q)^{1/p}}, \quad x \in (0,\infty), \tag{2.18}$$

then logarithmic differentiation leads to

$$\begin{aligned}\frac{F^{*\prime}(x)}{F^*(x)} =&(1-p_2)\frac{1}{x} - \frac{1}{\operatorname{arcsinh}_{p,q}(x)(1+x^q)^{1/p}} - \frac{qx^{q-1}}{p(1+x^q)}\\ =&\frac{1}{x}(1-p_2-G^*(x)),\end{aligned} \tag{2.19}$$

where

$$G^*(x) = \frac{x}{\operatorname{arcsinh}_{p,q}(x)(1+x^q)^{1/p}} + \frac{qx^q}{p(1+x^q)}.$$

Note that

$$\lim_{x\to 0} G^*(x) = 1. \tag{2.20}$$

It follows from Lemma 2.2 that $G^*(x) > 1$ for $x \in (0,\infty)$. Then we have $\inf\limits_{x\in(0,\infty)} G^*(x) = 1$. Hence equation (2.19) leads to the conclusion that $F^{*\prime}(x) \leq 0$ for $x \in (0,\infty)$ if and only if $p_2 \geq 0$. Namely, $F^*(x)$ is strictly decreasing on $(0,\infty)$ if and only if $p_2 \geq 0$. Hence if $p_2 \geq 0$, then $\partial J^*/\partial x < 0$ by (2.17), and $J^*(x,y) \geq J^*(y,y) = 1$. Then from (2.16) we have

$$\operatorname{arcsinh}_{p,q}(H_{p_2}(x,y)) \geq \sqrt{\operatorname{arcsinh}_{p,q}(x)\arcsin_{p,q}(y)} \tag{2.21}$$

for $p_2 \geq 0$, with equality if and only if $x = y$.

Taking $x = \sinh_{p,q}(r^*)$ and $y = \sinh_{p,q}(s^*)$ in inequality (2.21), we get

$$\sinh_{p,q}(\sqrt{r^*s^*}) \leq H_{p_2}(\sinh_{p,q}(r^*), \sinh_{p,q}(s^*)) \tag{2.22}$$

for all $r^*, s^* \in (0, m_{p,q})$ if $p_2 \geq 0$. In particular, inequality (1.2) follows from (2.22) with $p_2 = 0$.

Remark 2.1. Conjecture 1.1 follows easily from $\pi_{p,q}/2 > 1$ and Lemma 2.3 together with Theorem 1.1.

Acknowledgements

This research was supported by the Natural Science Foundation of China (Grant Nos. 11071059, 11071069, 11171307), and the Innovation Team Foundation of the Department of Education of Zhejiang Province (Grant No. T200924).

References

[1] G. D. Anderson, M. Vuorinen, and X.-H. Zhang, Topics on special function III, arXiv: 1209. 1696v1 [math. CA].

[2] B. A. Bhayo and M. Vuorinen, On generalized trigonometric functions with two parameters, J. Approx. Theory 164(2012), 1415-1426.

[3] B. A. Bhayo and M. Vuorinen, Inequalities for eigenfunctions of the p-Laplancian, ariXiv: 1101.3911v3 [math. CA].

[4] R. J. Biezuner, G. Ercole, E. M. Martins, Computing the first eigenvalue of the p-Laplacian via the inverse power method, J. Funct. Anal. 257(2009), 243-270.

[5] P. J. Bushell and D. E. Edmunds, Remarks on generalized trigonometric functions, Rocky Mountain J. Math. 42(2012), 25-57.

[6] P. Drábek, R. Manásevich, On the closed solution to some nonhomogenous eigenvalue problems with p-Laplacian, Differential Integral Equations 12(1999), 773-788.

[7] D. E. Edmunds, P. Gurka and J. Lang, Properties of generalized trigonometric functions, J. Approx. Theory 164(2012), 47-56.

[8] R. Klen, M. Vuorinen and X.-H. Zhang, Inequalities for the generalized trigonometric and hyperbolic functions, arXiv: 1210.6749v1 [math. CA].

[9] J. Lang, D. E. Edmunds, Eigenvalues, Embeddings and Generalized Trigonometric Functions, in: Lecture Notes in Mathematics 2016, Springer-Verlag, 2011.

[10] P. Lindqvist, Some remarkable sine and cosine functions, Ric. Mat. 44(1995), 269-290.

[11] P. Lindqvist and J. Peetre, p-arclength of the q-circle, Math. Student 72(2003), 139-145.

[12] S. Takeuchi, Generalized Jacobian elliptic functions and their applications to bifurcation problems associated with p-Laplacian, J. Math. Anal. Appl. 385(2012), 24-35.